



## Research

**Cite this article:** Boast AP, Wood JR, Cooper J, Bolstridge N, Perry GLW, Wilmshurst JM. 2025 DNA and spores from coprolites reveal that colourful truffle-like fungi endemic to New Zealand were consumed by extinct moa (*Dinornithiformes*). *Biol. Lett.* **21**: 20240440. <https://doi.org/10.1098/rsbl.2024.0440>

Received: 31 July 2024

Accepted: 25 November 2024

### Subject Category:

Global change biology

### Subject Areas:

palaeontology, ecology, ecosystem

### Keywords:

evolutionary anachronism, moa, New Zealand, extinction, ancient DNA, mycophagy

### Author for correspondence:

Alexander P. Boast

e-mail: [boasta@landcareresearch.co.nz](mailto:boasta@landcareresearch.co.nz)

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.7584142>.

# DNA and spores from coprolites reveal that colourful truffle-like fungi endemic to New Zealand were consumed by extinct moa (*Dinornithiformes*)

Alexander P. Boast<sup>1</sup>, Jamie R. Wood<sup>2,3</sup>, Jerry Cooper<sup>1</sup>, Nic Bolstridge<sup>1</sup>, George L. W. Perry<sup>4</sup> and Janet M. Wilmshurst<sup>1</sup>

<sup>1</sup>Manaaki Whenua-Landcare Research, Lincoln, Canterbury 7640, New Zealand

<sup>2</sup>School of Biological Sciences, Faculty of Science, Engineering and Technology, and <sup>3</sup>Environment Institute, University of Adelaide, Adelaide, South Australia 5005, Australia

<sup>4</sup>School of Environment, Faculty of Science, University of Auckland, Auckland 1010, New Zealand

**APB**, 0000-0001-6538-1316; **JRW**, 0000-0001-8008-6083; **JC**, 0000-0003-4994-3897; **NB**, 0000-0002-9206-3644; **GLWP**, 0000-0001-9672-9135; **JMW**, 0000-0002-4474-8569

Mycovores (animals that consume fungi) are important for fungal spore dispersal, including ectomycorrhizal (ECM) fungi symbiotic with forest-forming trees. As such, fungi and their symbionts may be impacted by mycophage extinction. New Zealand (NZ) has a diversity of unusual, colourful, endemic sequestrate (truffle-like) fungi, most of which are ECM. As NZ lacks native land mammals (except bats), and sequestrate fungi are typically drab and mammal-dispersed, NZ's sequestrate fungi are hypothesized to be adapted for bird dispersal. However, there is little direct evidence for this hypothesis, as 41% of NZ's native land bird species became extinct since initial human settlement in the thirteenth century. Here, we report ancient DNA and spores from the inside of two coprolites of NZ's extinct, endemic upland moa (*Megalapteryx didinus*) that reveal consumption and likely dispersal of ECM fungi, including at least one colourful sequestrate species. Contemporary data from NZ show that birds rarely consume fungi and that the introduced mammals preferentially consume exotic fungi. NZ's endemic sequestrate fungi could therefore be dispersal limited compared with fungi that co-evolved with mammalian dispersers. NZ's fungal communities may thus be undergoing a gradual species turnover following avian mycophage extinction and the establishment of mammalian mycophages, potentially affecting forest resilience and facilitating invasion by exotic tree taxa.

## 1. Introduction

Ecological networks often comprise thousands of interconnected species, meaning species losses can produce cascades that impact entire ecosystems [1]. Faunal declines and extinctions have characterized the late Quaternary (*ca* 50 ka to present), with megafauna and island endemics being among the worst affected [2–5]. The extent of recent broken species interactions may be reflected by the prevalence of taxa hypothetically adapted for absent partners (evolutionary anachronisms) [6,7]. For example, numerous plant species worldwide produce fruit that cannot be dispersed by indigenous extant species (e.g. owing to overlarge seeds or unpalatability) and were presumably dispersed by extinct fauna such as megafaunal mammals (e.g. proboscideans or ground sloths) [6–12] or giant tortoises [13–17]. Although some anachronistic plant taxa have benefitted from introduced surrogates [18–22], or have

been domesticated [23–25], others are range-restricted, recruitment/dispersal limited, or in severe decline [8,16,26–28], suggesting that the long-term effects of faunal loss are still unfolding in many ecosystems as ‘extinction debt’ [29–31].

Although disrupted animal–plant mutualisms are frequently presented as a likely consequence of faunal extinctions [32–37], how fungi have been affected by the loss of animal mutualists is poorly understood [38,39]. Animals that consume macrofungi sporocarps (mycophores) can disperse viable spores via their scats, resulting in an animal–fungi propagule dispersal mutualism [40–46]. Most spores shed by agaricoid (mushroom-forming) species tend to congregate near sporocarps, and those that successfully wind-disperse are highly diffuse in where they settle [47,48]. In contrast, animal-consumed spores are deposited in a rich growing medium at high concentrations and often far from the parent [43,49]. Furthermore, sequestrate (truffle-like) fungi do not forcibly discharge their spores, and mostly depend on mycophores for spore dispersal [50]. Sequestrate morphology has evolved independently across many macrofungal phyla and repeatedly within some genera (e.g. *Cortinarius* or *Russula*) [51–54], supporting that mycophores provide a reliable alternative for spore dispersal over wind transport. As many animal-dispersed fungi are ectomycorrhizal (ECM) with ecologically important plants (e.g. canopy-forming trees) [55], animal dispersal of ECM fungi can result in a tripartite animal–plant–fungal mutualism underpinning functions such as forest stability, productivity, regeneration and expansion [39,44,46,49,56–60]. Sequestrate fungal sporocarps are typically drab and often hypogeous (below-ground) and/or odorous [61–65], evidently reflecting adaptations for scent-foraging and digging species with poor colour vision (e.g. most mammals), rather than diurnal, visual-foraging species with low olfaction capability (e.g. most birds) [66]. Congruently, most known examples of vertebrate mycophores are mammals [67], with fewer documented examples of reptile [68] or bird [69,70] mycophagy.

Many sequestrate fungi endemic to New Zealand (NZ) produce brightly coloured, epigeous (above-ground), fruit-like sporocarps, which may be adaptations for bird dispersal [71–73]. However, it has been difficult to test the hypothesis that NZ’s fungi were bird-dispersed as most of NZ’s native bird species are either extinct or severely range-reduced. NZ’s long oceanic isolation from Gondwana (over 52 Ma) [74] produced a complex, mammal-free terrestrial biota (excluding bats or pre-Quaternary taxa) dominated by birds, including nine species of megaherbivore ratite moa (*Dinornithiformes*) spanning approximately 14–242 kg in weight [75,76]. The arrival of humans and introduced mammals to NZ after AD thirteenth century [77] resulted in the extinction of *ca* 41% of NZ’s native terrestrial bird species [78] and all moa by the sixteenth century [79,80]. Further, many of NZ’s surviving bird species are functionally extinct [81,82]. Avian mycophagy has rarely been observed in NZ, leading to speculations that NZ’s colourful sequestrate fungi are evolutionarily anachronistic and that the primary avian mycophores in NZ consisted of extirpated or extinct taxa such as moa [71,72,83]. NZ’s forests are heavily composed of widespread tree species co-dependent on ECM fungi, including southern beech (*Nothofagaceae*), mānuka (*Leptospermum scoparium*) and kānuka (*Kunzea ericoides*) [84]. As most of NZ’s sequestrate fungi are believed to be ECM [85], evidence for their dispersal by extinct birds would indicate that the full ecological consequences of these extinctions are yet to be realized in NZ forests [86].

Hundreds of Holocene avian coprolites, identified by ancient DNA (aDNA) as mostly being deposited by five species of moa and the endangered kākāpō parrot (*Strigops habroptila*), occur at several localities on NZ’s South Island [87]. Although a previous study on 18S rRNA metabarcoding of kākāpō, upland moa (*Megalapteryx didinus*), and South Island giant moa (*Dinornis robustus*) coprolites identified the presence of ECM fungi [86], taxonomic resolution was insufficient to confirm whether the fungi were sequestrate. Further, although moa coprolites have been previously analysed for palynomorphs [88–90] and parasites [91], they have not been investigated for fungal spore content and concentrations (spores mg<sup>−1</sup>). As a result, there is limited evidence that moa consumed sporocarps or truffle-like fungi specifically or played a role in the dispersal of their spores. Although there is circumstantial evidence that NZ’s sequestrate fungi are adapted for bird dispersal and were likely consumed by extinct birds such as moa [92], higher taxonomic resolution and spore evidence are needed to definitively answer the question: ‘did moa consume, and potentially disperse, NZ’s truffle-like fungi?’

## 2. Methods

Here, we report an opportunistic discovery from two coprolites that were originally sampled as part of a larger project on kākāpō coprolites and unexpectedly determined by aDNA as being from the relatively small (compared to other moa, approx. 14–63 kg) [76] upland moa [93]. The coprolites (figure 1) were from Hodges Creek (HC) Cave near the upper Tākaka River catchment in northwest Nelson (sample X/17/11/33, site elevation approx. 900 m), and Takahē Valley (TV) in Fiordland (sample X18/153/03, site elevation approx. 1100 m). HC Cave is surrounded by dense silver beech (*Lophozonia menziesii*) forest, and the coprolite (collected in 2018 by authors of this study) was located approximately 20 m into the cave system, partially covered in sediment and large rock fragments [93]. The TV sample, conversely, was from a museum specimen (CM 2016.171.11) with a less certain collection date and provenance. However, coprolite deposits in TV have been found in several rockshelters, and are all located adjacent to subalpine vegetation near to a silver beech-dominated treeline [93].

Specialist coprolite-subsampling procedures, <sup>14</sup>C dating, DNA extraction, DNA amplification and species identification of both samples are described in [93]. To minimize the risk of exogenous environmental contamination, the exterior of the coprolites was scraped off with a sterile scalpel, and the coprolite was then ultraviolet-radiated for 15 min before a subsample was taken from the interior of the coprolite under a sterile hood in a clean room designed for palaeoecological samples. All DNA extraction and PCR preparation were performed in a purpose-built, isolated aDNA laboratory. In addition to the two coprolite DNA extractions, we included extraction blank controls (EBCs, extractions performed without the addition of sample, to identify contaminants) from [93].


DNA extractions (both samples and EBCs) were amplified in triplicate for a region of the noncoding, nuclear gene Internal Transcribed Spacer 1 (ITS-1) using the fungi-specific primers ITS5 [94] and ITS5.8S [95] (initial 2 min denaturation of 94°C; 35





**Figure 1.** (a) Upland moa skeleton, (b) HC coprolite X17/11/33, and (c–h) examples of fungi identified from aDNA (\* denotes taxa with congruent spore evidence): (c) *Gallacea scleroderma*\*, (d) *Gallacea* sp. 'Nelson Lakes'\*, (e) *Rossbeevera pachydermis*\*, (f) *Russula macrocystidiata*\*, (g) *Cortinarius* sp. 'Blyth Track', (h) *Cortinarius violaceovolvatus*. Photo credits: (a) Wikimedia Commons, (b) Alexander P. Boast, (c–h) Noah Siegel.

cycles of 30 s at 94°C, 30 s at 60°C and 40 s at 72°C; and a final extension of 10 min at 72°C). We selected ITS5 and ITS5.8 as these primers have been shown to reliably amplify a broad diversity of fungi (including Ascomycota and Basidiomycota) from ancient samples [95]. Reagents, PCR preparation, indexing, quantification, library preparation and sequencing of amplicons, as well as early bioinformatic processing (e.g. demultiplexing, primer removal and quality filtering) followed [93]. Identifications used a reference database built from all ITS fungal sequences on GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) identified to family level. FASTA files representing all amplicon sequence variants (ASVs) were analysed using the BLASTn algorithm in BLAST+ v. 2.7.1 [96], against our database, using parameters described in [93]. BLASTn outputs were processed using MEGAN v. 6 [97], using parameters optimized to detect taxa rather than provide conservative identifications: the top 5% of hits, a minimum percentage identity of 80%, and a bitscore cut-off of 200. All ASVs identified in an EBC control were removed from subsequent analyses. Further, all ASVs attributed to macrofungi or ECM fungi (none of which appeared in EBC controls) had their taxonomic identity confirmed by maximum-clade credibility Bayesian phylogenetic analyses in MrBayes, using a series of selected outgroup taxa and a substitution model identified by jModeltest v. 2.1.10 [98],  $1 \times 10^7$  generations and discarding 25% of trees as a burn-in (electronic supplementary material, figures S1–S6).



**Figure 2.** Percentage of reads assigned to taxa from the two coprolite samples, subdivided into all fungi (left) and Agaricomycetes only (right). Taxa coloured blue are those with congruent spore evidence in the coprolites, and most likely to be dietary.

Spore identification was performed on microfossil slides prepared using methods in [93], omitting the acetolysis and staining steps. During microfossil preparation, a known number of exotic *Lycopodium clavatum* marker spores were added (University of Lund, batch no. 1031, 20 848 spores per tablet, on average). Fungal spores were identified at 1000× magnification (oil immersion) and were compared with those available for collections deposited in the Plant Disease Division Fungarium, using online imagery available via the Systematic Collection Data website (<https://scd.landcareresearch.co.nz/>) and the Biota of New Zealand website (<https://biotanz.landcareresearch.co.nz/>). Fungal spores and *L. clavatum* spores were subsequently counted at 200× magnification until at least 250 fungal spores were counted. Known weights of the microfossil subsamples, compared with *L. clavatum* counts, meant that the mass of counted samples and fungal spore concentration could be estimated.

### 3. Results

Fungal DNA assemblages from both HC and TV [99] were dominated by saprobic, microfungal Mucoromycota and Ascomycota (figure 2; electronic supplementary material, dataset 1), which could not be confidently distinguished from common environmental species, were unlikely to be dietary, and were not analysed further. All remaining fungal diversity comprised macrofungal Basidiomycota in the Agaricomycetes (6.4 and 25.5% of reads from the HC and TV samples, respectively; figure 2). Unlike microfungal Ascomycota or Mucoromycota, macrofungal Agaricomycetes had the potential to be dietary. For example, all genera of Agaricomycetes we detect produce sporocarps large enough to be foraged on by mycophages (electronic supplementary material, dataset 1). Further, most identified Agaricomycetes sequences were of habitat-specific ECM taxa, which are unlikely to be laboratory contaminants (e.g. no ECM fungi were detected in EBC controls) or to colonize coprolites *in situ* (neither coprolite was associated with forest soil, and ECM fungi depend on symbiotic tree roots for development) [86]. Hereafter, percentage statistics are derived from Agaricomycetes only.

Thirteen ASVs were identified as native macrofungal Agaricomycetes. Most were identifiable to species, either by close pairwise match (nine showed more than 98.5% similarity to reference sequences) (electronic supplementary material, dataset 1) and/or by phylogenetic association (electronic supplementary material, figures S1–S6). Of these, ECM sequestered fungi were dominant (figure 1), especially the purple *Gallaceae scleroderma* (95.5% of HC and 14.8% of TV), the purple *Cortinarius violaceovolvatus* (7.4% of TV) and the drab (but staining blue with bruising or exposure to air) *Rosbeevera pachydermis* (7.4% of TV). Low proportions of HC reads were also identified as falling within, or related to, a clade of two near-identical, drab/pale pink *Gallaceae* species (*G. eburnea*, *G. sp.* 'Nelson Lakes'). Agaricoid species included the ECM *Russula macrocystidiata*, and *Cortinarius* spp., the saprobic *Armillaria novae-zelandiae* and an unidentified *Pholiota* (a saprobic genus). High concentrations of spores with morphologies congruent with those of the ECM fungi genera *Gallaceae* were identified from HC (1885 spores mg<sup>-1</sup>, occasionally in clusters admixed with fungal tissue) and *Rosbeevera* and *Russula* from TV (7205 and 3556 spores mg<sup>-1</sup>, respectively) (electronic supplementary material, figures S7 and S8). These spore concentrations fall within the range observed in the dung (by dry weight) of contemporary avian mycophages, and strongly support sporocarp consumption [70].



## 4. Discussion

Here, for the first time, we present co-occurring fungal spore and species-level fungal DNA evidence from moa coprolites. Together, these two sources of data provide definitive support that upland moa consumed the sporocarps of ECM fungi, including brightly coloured sequestrate species (figures 1 and 2). ECM fungi are extremely unlikely to colonize coprolites *in situ*, as ECM fungi depend on symbiotic tree roots for development and would therefore have been unable to colonize moa coprolites located in dry, root-free sediment away from forest soil. Furthermore, those fungal species with supporting DNA and spore evidence were low in richness (one sp. HC, two spp. TV) and were represented by high spore concentrations. In comparison, environmental spore contamination or incidental spore consumption by moa would likely result in a wider variety of spore types and much lower spore concentrations. Further, consumption of other fungal tissues is not supported (e.g. moa incidentally consuming hyphae and forest soil when feeding on the forest floor), as such tissues are not associated with spores. Conversely, our spore data would have been of limited value without DNA evidence, as fungal spores are extremely difficult to identify to species through morphology alone. Our study thus builds considerably on the study of Boast *et al.* [86], who identified ECM-specific fungal genera from moa coprolites (*Cortinarius*, *Inocybe*) but lacked spore evidence and were unable to provide species-level resolution or detect sequestrate species. Overall, this new discovery confirms an interaction between two geographically separated moa (almost 600 km) and coloured sequestrate fungi (and ECM fungi in general). As consumption is also likely to have resulted in spore dispersal, we thus infer that this interaction represents a lost mutualism.

At least three species of ECM fungi had co-occurring spore and DNA observations in our data: at least one species of the sequestrate genus *Gallaceae* (strongly supported by DNA read percentages to be the purple *G. scleroderma*), the drab to blue sequestrate species *R. pachydermis* and the agaricoid species *R. macrocystidiata* (figure 1). Although other fungi detected in the DNA data were not observed in the spore counts (e.g. the ‘purple pouch fungus’ *C. violaceovolatus*), these could represent consumed sporocarps in some cases. For example, if sporocarps were consumed prior to sporulation, spores would either be absent or potentially too undeveloped to survive digestion, preservation or laboratory procedures [100]. All moa species with gizzard or coprolite records consumed brightly coloured fleshy fruit, suggesting that they were visual foragers and would have been attracted to similarly coloured fungi [101]. Other ratites, such as emu (*Dromaius novaehollandiae*) [102] and southern cassowary (*Casuaris casuaris*) [103] consume agaricoid fungi, and upland moa, giant moa (*Dinornis* spp.) and little bush moa (*Anomalopteryx didiformis*) foraged on forest floors [86,88,90,101,104]. While our data demonstrate that at least one moa species consumed sequestrate fungi, it is likely that most other moa species did too, given their generalist diets and high degree of dietary overlap between species [86,88,101]. Further analyses of moa coprolites will be required to reveal whether sequestrate fungi consumption was a common and widespread activity among moa.

Although NZ's presettlement avifauna was diverse, moa may have been exceptional fungal dispersers. The fibrous diets of herbivorous moa, osteological evidence and allometric scaling imply that moa had larger digestive systems and home ranges and longer gut retention times than other NZ ground-dwelling birds [101,105–107]. An extant large ratite, the common ostrich (*Struthio camelus*), has an average retention time for small particulates of 30–36 h [108]. Moa could thus have dispersed spores from any consumed fungus across substantial distances and during several deposition events. Some moa species had broad ranges, with upland moa evidently moving between lowlands, forests and alpine herbfields [75,90,109], meaning they could have contributed to the fungal spore inoculation of soils in non-forest habitats and thereby assisted with forest spread and regeneration [57,60]. Of NZ's surviving avifauna, only the critically endangered kākāpō, North Island robin (*Petroica longipes*) and weka (*Gallirallus australis*) have been observed to consume fungi [69,73]. However, robins and other small birds (e.g. passerines, including introduced species) may be limited dispersers, as they have short gut retention times and typically have small home ranges [110–112]. Most larger herbivorous or omnivorous ground-dwelling birds in NZ are extinct [75] or endangered [82], with only the omnivorous rail weka locally common in some regions of NZ [113]. Few introduced ground-dwelling birds occur deep within native forests [114]. Overall, moa have no functional analogues as fungal dispersers among NZ's contemporary avifauna, and the overall contribution of birds to fungal dispersal in NZ must now be heavily diminished relative to presettlement NZ.

Coloured and non-scented sequestrate fungi appear to be poorly adapted for mammal dispersal relative to bird dispersal. Brightly coloured sequestrate fungi also occur in New Caledonia (e.g. *Leratiomyces similis*, *L. smagdarinus*) [115] and Lord Howe Island (*Secotium fragariosum*) [116] which, like NZ, lack indigenous mammals (except bats) and had vertebrate faunas dominated by birds and/or reptiles [92,117,118]. Further, compared with NZ, a very low proportion of sequestrate fungi on continental ecosystems are brightly coloured (e.g. *Cortinarius atratus* in southern Australia, *Halligena purpurea* in Patagonia) [70,119]. In NZ, introduced red deer (*Cervus elaphus*) and Australian brushtail possum (*Trichosurus vulpecula*) were observed consuming both native and exotic fungi, although only exotic fungal spores were viable post-digestion [120]. Further, in that same study [120], mammals preferentially foraged on sequestrate species that were exotic (*Rhizopogon*) or shared with Australia (*Octaviania*) [121]. As introduced mammals have largely replaced NZ's avian-dominated ground vertebrates, it can be assumed that sporocarps attractive to mammals now have an adaptive advantage over many native fungi species, being more likely to be foraged on and/or successfully dispersed.

A guild of mycophores in forested ecosystems results in a more diverse and admixed fungal community, benefitting forest productivity, regeneration and resilience [39]. Animals can inoculate unforested areas with ECM fungal spores, which may assist with forest spread or regeneration [57]. For example, the absence of symbiotic ECM in soil can limit the establishment of NZ beech [122,123]. The removal of native fungi dispersers, such as moa, from NZ's ecosystems has therefore likely resulted in a reduced dispersal capacity for many NZ fungi, possibly reducing rates of spread and regeneration of some forest types as a result. Some of NZ's ECM fungi may now be developing increasingly clustered distributions at the local scale, characteristic of dispersal-limited taxa [39], with flow-on effects for forest resilience. Further, a suite of species attractive to mammalian

dispersers may now dominate mycove faecal ECM inoculum in NZ (e.g. in unforested areas), conceivably causing NZ's ECM communities to become increasingly biased towards non-endemic taxa [120]. This may further impact forest ecology by opening invasion pathways for trees better adapted for symbioses with exotic fungi (e.g. lodgepole pine, *Pinus contorta*) [120,124–126].

We provide robust evidence, using coprolites, that native NZ fungi were consumed by the extinct moa. Further, we strongly support the hypothesis that many of NZ's ECM native fungi were adapted for, or at least benefitted from, avian dispersal, an interaction likely to have been drastically reduced by NZ bird extinction and range contraction. Our small sample size (two coprolites) provides a key starting point for identifying this novel interaction. However, further analysis of more coprolites and modern scats from a range of different species will help to address the question of how bird–fungi interactions in NZ forests have changed over time. In conclusion, we argue that NZ's sequestrate fungi present a strong case for evolutionary anachronism, and the long-term consequences of the lost native mycove, such as moa, may yet be fully realized in NZ's forests.

**Ethics.** This work did not require ethical approval from a human subject or animal welfare committee.

**Data accessibility.** Supporting data for this study—including raw and processed metabarcoding DNA data, alignments for phylogenetic analyses and custom-built reference database—are available on the Manaaki Whenua–Landcare Research Datastore [99].

Supplementary material is available online [127].

**Declaration of AI use.** We have not used AI-assisted technologies in creating this article.

**Authors' contributions.** A.P.B.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, software, visualization, writing—original draft, writing—review and editing; J.R.W.: conceptualization, methodology, project administration, resources, software, supervision, validation, visualization, writing—review and editing; J.C.: investigation, resources, validation, writing—review and editing; N.B.: investigation, methodology, project administration, resources, software, supervision, writing—review and editing; G.L.W.P.: conceptualization, funding acquisition, methodology, project administration, supervision, writing—review and editing; J.M.W.: conceptualization, funding acquisition, methodology, project administration, resources, software, supervision, visualization, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

**Conflict of interest declaration.** We declare we have no competing interests.

**Funding.** This project was funded by the University of Auckland, the Ornithological Society of New Zealand, the Royal Society of New Zealand, the Linnean Society of New South Wales and Strategic Science Investment Funding for Crown Research Institutes from the New Zealand Ministry of Business, Innovation and Employment's Science and Innovation Group.

**Acknowledgements.** We wish to thank Canterbury Museum for loaning specimens; Trevor Worthy and Paul Scofield for advice on sampling; John Patterson, Michal Kuchar and Nia-Joelle Weinzwieg for field support. We also wish to acknowledge and thank the indigenous kaitiaki (guardians) of the Tākaka River system (Manawhenua ki Mohua and Ngāti Apa ki te Rā Tō), and of Takahē Valley (Kāi Tahu Ōraka Aparima).

## References

- Montoya JM, Pimm SL, Solé RV. 2006 Ecological networks and their fragility. *Nature* **442**, 259–264. (doi:10.1038/nature04927)
- Martin PS, Steadman DW. 1999 Prehistoric extinctions on islands and continents. In *Extinctions in near time* (ed. RDE MacPhee), pp. 17–55. Boston, MA: Springer. (doi:10.1007/978-1-4757-5202-1\_2)
- Ripple WJ *et al.* 2015 Collapse of the world's largest herbivores. *Sci. Adv.* **1**, e1400103. (doi:10.1126/sciadv.1400103)
- Steadman DW. 1995 Prehistoric extinctions of Pacific island birds: biodiversity meets zooarchaeology. *Science* **267**, 1123–1131. (doi:10.1126/science.267.5201.1123)
- Wolf C, Ripple WJ. 2017 Range contractions of the world's large carnivores. *R. Soc. Open Sci.* **4**, 170052. (doi:10.1098/rsos.170052)
- Barlow C. 2008 *The ghosts of evolution: nonsensical fruit, missing partners, and other ecological anachronisms*. New York, NY: Basic Books.
- Janzen DH, Martin PS. 1982 Neotropical anachronisms: the fruits the gomphotheres ate. *Science* **215**, 19–27. (doi:10.1126/science.215.4528.19)
- Godfrey LR, Jungers WL, Schwartz GT, Irwin MT. 2008 Ghosts and orphans: Madagascar's vanishing ecosystems. In *Elwyn Simons: a search for origins*, pp. 361–395. New York, NY: Springer. (doi:10.1007/978-0-387-73896-3\_24)
- Crowley BE, Godfrey LR, Irwin MT. 2011 A glance to the past: subfossils, stable isotopes, seed dispersal, and lemur species loss in southern Madagascar. *Am. J. Primatol.* **73**, 25–37. (doi:10.1002/ajp.20817)
- Galetti M *et al.* 2018 Ecological and evolutionary legacy of megafauna extinctions. *Biol. Rev.* **93**, 845–862. (doi:10.1111/brv.12374)
- Newsom LA, Muhlbachler MC. 2006 Mastodons (*Mammot americanum*) diet foraging patterns based on analysis of dung deposits. In *First floridians and last mastodons: the Page-Ladson site in the Aucilla River*, pp. 263–331. Dordrecht, The Netherlands: Springer. (doi:10.1007/978-1-4020-4694-0\_10)
- Barlow CC. 2001 Anachronistic fruits and the ghosts who haunt them. *Arnoldia* **61**, 14–21. (doi:10.5962/p.390407)
- Steadman DW, Albury NA, Carlson LA, Franz R, LeFebvre MJ, Kakuk B, Keegan WF. 2020 The paleoecology and extinction of endemic tortoises in the Bahamian Archipelago. *Holocene* **30**, 420–427. (doi:10.1177/0959683619887412)
- Andriantsaralaza S, Pedrono M, Tassin J, Roger E, Rakouth B, Danthu P. 2014 The role of extinct giant tortoises in the germination of extant baobab *Adansonia rubrostipa* seeds in Madagascar. *Afr. J. Ecol.* **52**, 246–249. (doi:10.1111/aje.12101)
- Moolna A. 2008 Preliminary observations indicate that giant tortoise ingestion improves seed germination for an endemic ebony species in Mauritius. *Afr. J. Ecol.* **46**, 217–219. (doi:10.1111/j.1365-2028.2007.00807.x)
- Albert-Daviaud A *et al.* 2020 The ghost fruits of Madagascar: identifying dysfunctional seed dispersal in Madagascar's endemic flora. *Biol. Conserv.* **242**, 108438. (doi:10.1016/j.biocon.2020.108438)
- Hansen DM, Donlan CJ, Griffiths CJ, Campbell KJ. 2010 Ecological history and latent conservation potential: large and giant tortoises as a model for taxon substitutions. *Ecography* **33**, 272–284. (doi:10.1111/j.1600-0587.2010.06305.x)
- Brown JR, Archer S. 1989 Woody plant invasion of grasslands: establishment of honey mesquite (*Prosopis glandulosa* var. *glandulosa*) on sites differing in herbaceous biomass and grazing history. *Oecologia* **80**, 19–26. (doi:10.1007/BF00789926)



19. Janzen DH. 1982 Fruit traits, and seed consumption by rodents, of *Crescentia alata* (Bignoniaceae) in Santa Rosa National Park, Costa Rica. *Am. J. Bot.* **69**, 1258–1268. (doi:10.1002/j.1537-2197.1982.tb13371.x)
20. Janzen DH. 1982 Differential seed survival and passage rates in cows and horses, surrogate Pleistocene dispersal agents. *Oikos* **38**, 150–156. (doi:10.2307/3544014)
21. Janzen DH. 1981 Enterolobium cyclocarpum seed passage rate and survival in horses, Costa Rican Pleistocene seed dispersal agents. *Ecology* **62**, 593–601. (doi:10.2307/1937726)
22. Janzen DH. 1981 Guanacaste tree seed-swallowing by Costa Rican range horses. *Ecology* **62**, 587–592. (doi:10.2307/1937725)
23. Kistler L, Newsom LA, Ryan TM, Clarke AC, Smith BD, Perry GH. 2015 Gourds and squashes (*Cucurbita* spp.) adapted to megafaunal extinction and ecological anachronism through domestication. *Proc. Natl Acad. Sci. USA* **112**, 15107–15112. (doi:10.1073/pnas.1516109112)
24. Galindo-Tovar ME, Ogata-Aguilar N, Arzate-Fernández AM. 2008 Some aspects of avocado (*Persea americana* Mill.) diversity and domestication in Mesoamerica. *Genet. Resour. Crop Evol.* **55**, 441–450. (doi:10.1007/s10722-007-9250-5)
25. Thomas E, van Zonneveld M, Loo J, Hodgkin T, Galluzzi G, van Etten J. 2012 Present spatial diversity patterns of *Theobroma cacao* L. in the neotropics reflect genetic differentiation in Pleistocene refugia followed by human-influenced dispersal. *PLoS ONE* **7**, e47676. (doi:10.1371/journal.pone.0047676)
26. Jackson PSW, Cronk QC, Parnell JA. 1988 Notes on the regeneration of two rare Mauritian endemic trees. *Trop. Ecol.* **78**, 56–65.
27. Hall JA, Walter GH. 2013 Seed dispersal of the Australian cycad *Macrozamia miquelii* (Zamiaceae): are cycads megafauna-dispersed 'grove forming' plants? *Am. J. Bot.* **100**, 1127–1136. (doi:10.3732/ajb.1200115)
28. Zaya DN, Howe HF. 2009 The anomalous Kentucky coffeetree: megafaunal fruit sinking to extinction? *Oecologia* **161**, 221–226. (doi:10.1007/s00442-009-1372-3)
29. Colwell RK, Dunn RR, Harris NC. 2012 Coextinction and persistence of dependent species in a changing world. *Annu. Rev. Ecol. Evol. Syst.* **43**, 183–203. (doi:10.1146/annurev-ecolsys-110411-160304)
30. Dunn RR, Harris NC, Colwell RK, Koh LP, Sodhi NS. 2009 The sixth mass coextinction: are most endangered species parasites and mutualists? *Proc. R. Soc. B* **276**, 3037–3045. (doi:10.1098/rspb.2009.0413)
31. Blanchard G, Munoz F. 2023 Revisiting extinction debt through the lens of multitrophic networks and meta-ecosystems. *Oikos* **2023**, e09435. (doi:10.1111/oik.09435)
32. Pires MM, Guimarães PR, Galetti M, Jordano P. 2018 Pleistocene megafaunal extinctions and the functional loss of long-distance seed-dispersal services. *Ecography* **41**, 153–163. (doi:10.1111/ecog.03163)
33. Johnson CN. 2009 Ecological consequences of Late Quaternary extinctions of megafauna. *Proc. R. Soc. B* **276**, 2509–2519. (doi:10.1098/rspb.2008.1921)
34. Heinen JH, van Loon EE, Hansen DM, Kissling WD. 2018 Extinction-driven changes in frugivore communities on oceanic islands. *Ecography* **41**, 1245–1255. (doi:10.1111/ecog.03462)
35. McConkey KR, Drake DR. 2006 Flying foxes cease to function as seed dispersers long before they become rare. *Ecology* **87**, 271–276. (doi:10.1890/05-0386)
36. McConkey KR, O'Farrill G. 2016 Loss of seed dispersal before the loss of seed dispersers. *Biol. Conserv.* **201**, 38–49. (doi:10.1016/j.biocon.2016.06.024)
37. Fricke EC, Tewksbury JJ, Rogers HS. 2018 Defaunation leads to interaction deficits, not interaction compensation, in an island seed dispersal network. *Glob. Chang. Biol.* **24**, e190–e200. (doi:10.1111/gcb.13832)
38. Nuske SJ, Vernes K, May TW, Claridge AW, Congdon BC, Krockenberger A, Abell SE. 2017 Redundancy among mammalian fungal dispersers and the importance of declining specialists. *Fungal Ecol.* **27**, 1–13. (doi:10.1016/j.funeco.2017.02.005)
39. Nuske SJ, Anslan S, Tedersoo L, Congdon BC, Abell SE. 2019 Ectomycorrhizal fungal communities are dominated by mammalian dispersed truffle-like taxa in north-east Australian woodlands. *Mycorrhiza* **29**, 181–193. (doi:10.1007/s00572-019-00886-2)
40. Colgan W III, Claridge AW. 2002 Mycorrhizal effectiveness of *Rhizopogon* spores recovered from faecal pellets of small forest-dwelling mammals. *Mycol. Res.* **106**, 314–320. (doi:10.1017/s0953756202005634)
41. Trappe JM, Maser C. 1976 Germination of spores of *Glomus macrocarpum* (Endogonaceae) after passage through a rodent digestive tract. *Mycologia* **68**, 433–436. (doi:10.2307/3759016)
42. Cork SJ, Kenagy GJ. 1989 Rates of gut passage and retention of hypogeous fungal spores in two forest-dwelling rodents. *J. Mammal.* **70**, 512–519. (doi:10.2307/1381423)
43. Claridge AW, Tanton MT, Seebeck JH, Cork SJ, Cunningham RB. 1992 Establishment of ectomycorrhizae on the roots of two species of *Eucalyptus* from fungal spores contained in the faeces of the long-nosed potoroo (*Potorous tridactylus*). *Aust. J. Ecol.* **17**, 207–217. (doi:10.1111/j.1442-9993.1992.tb00799.x)
44. Maser C, Trappe JM, Nussbaum RA. 1978 Fungal–small mammal interrelationships with emphasis on Oregon coniferous forests. *Ecology* **59**, 799–809. (doi:10.2307/1938784)
45. Kotter MM, Farentinos RC. 1984 Formation of Ponderosa pine ectomycorrhizae after inoculation with feces of tassel-eared squirrels. *Mycologia* **76**, 758–760. (doi:10.1080/00275514.1984.12023910)
46. Lamont BB, Ralph CS, Christensen PES. 1985 Mycophagous marsupials as dispersal agents for ectomycorrhizal fungi on *Eucalyptus calophylla* and *Gastrolobium bilobum*. *New Phytol.* **101**, 651–656. (doi:10.1111/j.1469-8137.1985.tb02870.x)
47. Galante TE, Horton TR, Swaney DP. 2011 95% of basidiospores fall within 1 m of the cap: a field- and modeling-based study. *Mycologia* **103**, 1175–1183. (doi:10.3852/10-388)
48. Li DW. 2005 Release and dispersal of basidiospores from *Amanita muscaria* var. *alba* and their infiltration into a residence. *Mycol. Res.* **109**, 1235–1242. (doi:10.1017/s0953756205003953)
49. Johnson CN. 1996 Interactions between mammals and ectomycorrhizal fungi. *Trends Ecol. Evol.* **11**, 503–507. (doi:10.1016/s0169-5347(96)10053-7)
50. Thiers HD. 1984 The secotoid syndrome. *Mycologia* **76**, 1–8. (doi:10.1080/00275514.1984.12023803)
51. Peintner U, Bougher NL, Castellano MA, Moncalvo J, Moser MM, Trappe JM, Vilgalys R. 2001 Multiple origins of sequestrate fungi related to *Cortinarius* (Cortinariaceae). *Am. J. Bot.* **88**, 2168–2179. (doi:10.2307/3558378)
52. Bruns TD, Fogel R, White TJ, Palmer JD. 1989 Accelerated evolution of a false-truffle from a mushroom ancestor. *Nature* **339**, 140–142. (doi:10.1038/339140a0)
53. Miller SL, McClean TM, Walker JF, Buyck B. 2001 A molecular phylogeny of the Russulales including agaricoid, gasteroid and pleurotoid taxa. *Mycologia* **93**, 344–354. (doi:10.1080/00275514.2001.12063166)
54. Wilson AW, Binder M, Hobbitt DS. 2011 Effects of gasteroid fruiting body morphology on diversification rates in three independent clades of fungi estimated using binary state speciation and extinction analysis. *Evolution* **65**, 1305–1322. (doi:10.1111/j.1558-5646.2010.01214.x)
55. Marks GC, Kozłowski TT. 1973 *Ectomycorrhizae, their ecology and physiology*. New York, NY: Academic Press.
56. Malajczuk N, Trappe JM, Molina R. 1987 Interrelationships among some ectomycorrhizal trees, hypogeous fungi and small mammals: Western Australian and northwestern American parallels. *Aust. J. Ecol.* **12**, 53–55. (doi:10.1111/j.1442-9993.1987.tb00927.x)
57. Cázares E, Trappe JM. 1994 Spore dispersal of ectomycorrhizal fungi on a glacier forefront by mammal mycophagy. *Mycologia* **86**, 507–510. (doi:10.1080/00275514.1994.12026443)
58. Terwilliger J, Pastor J. 1999 Small mammals, ectomycorrhizae, and conifer succession in beaver meadows. *Oikos* **85**, 83–94. (doi:10.2307/3546794)

59. Johnson CN. 1995 Interactions between fire, mycophagous mammals, and dispersal of ectomycorrhizal fungi in eucalyptus forests. *Oecologia* **104**, 467–475. (doi:10.1007/BF00341344)
60. Frank JL, Anglin S, Carrington EM, Taylor DS, Viratos B, Southworth D. 2009 Rodent dispersal of fungal spores promotes seedling establishment away from mycorrhizal networks on *Quercus garryana*. *Botany* **87**, 821–829. (doi:10.1139/b09-044)
61. Splivallo R, Ottonello S, Mello A, Karlovsky P. 2011 Truffle volatiles: from chemical ecology to aroma biosynthesis. *New Phytol.* **189**, 688–699. (doi:10.1111/j.1469-8137.2010.03523.x)
62. Rapior S, Mauruc MJ, Guinberteau J, Masson CL, Bessière JM. 2000 Volatile composition of *Gyrophragmium dunali*. *Mycologia* **92**, 1043–1046. (doi:10.1080/00275514.2000.12061249)
63. Donaldson R, Stoddart M. 1994 Detection of hypogeous fungi by Tasmanian bettong (*Bettongia gaimardi*: Marsupialia; Macropodoidea). *J. Chem. Ecol.* **20**, 1201–1207. (doi:10.1007/BF02059754)
64. Talou T, Gaset A, Delmas M, Kulifaj M, Montant C. 1990 Dimethyl sulphide: the secret for black truffle hunting by animals? *Mycol. Res.* **94**, 277–278. (doi:10.1016/s0953-7562(09)80630-8)
65. Stephens RB, Trowbridge AM, Ouimette AP, Knighton WB, Hobbie EA, Stoy PC, Rowe RJ. 2020 Signaling from below: rodents select for deeper fruiting truffles with stronger volatile emissions. *Ecology* **101**, e02964. (doi:10.1002/ecy.2964)
66. Corfield JR, Price K, Iwaniuk AN, Gutierrez-Ibañez C, Birkhead T, Wylie DR. 2015 Diversity in olfactory bulb size in birds reflects allometry, ecology, and phylogeny. *Front. Neuroanat.* **9**, 102. (doi:10.3389/fnana.2015.00102)
67. Elliott TF, Truong C, Jackson SM, Zúñiga CL, Trappe JM, Vernes K. 2022 Mammalian mycophagy: a global review of ecosystem interactions between mammals and fungi. *Fungal Syst. Evol.* **9**, 99–159. (doi:10.3114/fuse.2022.09.07)
68. Elliott T, Bower D, Vernes K. 2019 Reptilian mycophagy: a global review of mutually beneficial associations between reptiles and macrofungi. *Mycosphere* **10**, 776–797. (doi:10.5943/mycosphere/10/1/18)
69. Elliott TF, Jusino MA, Trappe JM, Lepp H, Ballard GA, Bruhl JJ, Vernes K. 2019 A global review of the ecological significance of symbiotic associations between birds and fungi. *Fungal Divers.* **98**, 161–194. (doi:10.1007/s13225-019-00436-3)
70. Caiafa MV, Jusino MA, Wilkie AC, Díaz IA, Sieving KE, Smith ME. 2021 Discovering the role of Patagonian birds in the dispersal of truffles and other mycorrhizal fungi. *Curr. Biol.* **31**, 5558–5570. (doi:10.1016/j.cub.2021.10.024)
71. Beever RE. 1993 Dispersal of truffle-like fungi in New Zealand. In *Southern temperate ecosystems: origin and diversification* (ed. RS Hill), p. 439. Hobart, Australia: Australian Systematic Botany Society.
72. Beever R, Lebel T. 2014 Truffles of New Zealand: a discussion of bird dispersal characteristics of fruit bodies. *Auck. Bot. J.* **6**, 170–177.
73. Johns GB. 2022 Dispersal and trait evolution: did New Zealand's sequester fungi evolve unique traits to match bird sensory ecologies? MSc thesis, University of Auckland, New Zealand.
74. Schellart WP, Lister GS, Toy VG. 2006 A Late Cretaceous and Cenozoic reconstruction of the southwest Pacific region: tectonics controlled by subduction and slab rollback processes. *Earth Sci. Rev.* **76**, 191–233. (doi:10.1016/j.earscirev.2006.01.002)
75. Worthy TH, Holdaway RN. 2002 *The lost world of the moa: prehistoric life of New Zealand*. Christchurch, New Zealand: Canterbury University Press.
76. Tennyson AJD, Martinson P. 2006 *Extinct birds of New Zealand*. Wellington, New Zealand: Te Papa Press.
77. Wilmshurst JM, Anderson AJ, Higham TFG, Worthy TH. 2008 Dating the late prehistoric dispersal of Polynesians to New Zealand using the commensal Pacific rat. *Proc. Natl Acad. Sci. USA* **105**, 7676–7680. (doi:10.1073/pnas.0801507105)
78. Wood JR. 2023 Post-settlement extinction rates for the New Zealand avifauna. *N. Z. J. Ecol.* **47**, 3506. (doi:10.20417/nzjecol.47.3506)
79. Perry GLW, Wheeler AB, Wood JR, Wilmshurst JM. 2014 A high-precision chronology for the rapid extinction of New Zealand moa (Aves, Dinornithiformes). *Quat. Sci. Rev.* **105**, 126–135. (doi:10.1016/j.quascirev.2014.09.025)
80. Tomlinson S *et al.* 2024 Ecological dynamics of moa extinctions reveal convergent refugia that today harbour flightless birds. *Nat. Ecol. Evol.* **8**, 1472–1481. (doi:10.1038/s41559-024-02449-x)
81. Kelly D, Ladley JJ, Robertson AW, Anderson SH, Wotton DM, Wiser SK. 2010 Mutualisms with the wreckage of an avifauna: the status of bird pollination and fruit-dispersal in New Zealand. *N. Z. J. Ecol.* **34**, 66–85.
82. Robertson HA. 2021 *Conservation status of birds in Aotearoa New Zealand, 2021*. Wellington, New Zealand: Department of Conservation Te Papa Atawhai.
83. Beever R. 1999 Dispersal of New Zealand sequester fungi. In *IUMS 9th Int. Congress of Mycology, Sydney, Australia, 16–20 August 1999*, p. 190.
84. Orlovich DA, Cairney JohnWG. 2004 Ectomycorrhizal fungi in New Zealand: current perspectives and future directions. *N. Z. J. Bot.* **42**, 721–738. (doi:10.1080/0028825x.2004.9512926)
85. Bougher NL, Lebel T. 2001 Sequester (truffle-like) fungi of Australia and New Zealand. *Aust. Syst. Bot.* **14**, 439484. (doi:10.1071/sb00002)
86. Boast AP, Weyrich LS, Wood JR, Metcalf JL, Knight R, Cooper A. 2018 Coprolites reveal ecological interactions lost with the extinction of New Zealand birds. *Proc. Natl Acad. Sci. USA* **115**, 1546–1551. (doi:10.1073/pnas.1712337115)
87. Wood JR, Wilmshurst JM. 2014 Late Quaternary terrestrial vertebrate coprolites from New Zealand. *Quat. Sci. Rev.* **98**, 33–44. (doi:10.1016/j.quascirev.2014.05.020)
88. Wood JR, Wilmshurst JM, Richardson SJ, Rawlence NJ, Wagstaff SJ, Worthy TH, Cooper A. 2013 Resolving lost herbivore community structure using coprolites of four sympatric moa species (Aves: Dinornithiformes). *Proc. Natl Acad. Sci. USA* **110**, 16910–16915. (doi:10.1073/pnas.1307700110)
89. Wood J, Rawlence N, Rogers G, Austin J, Worthy T, Cooper A. 2008 Coprolite deposits reveal the diet and ecology of the extinct New Zealand megaherbivore moa (Aves, Dinornithiformes). *Quat. Sci. Rev.* **27**, 2593–2602. (doi:10.1016/j.quascirev.2008.09.019)
90. Wood JR, Wilmshurst JM, Wagstaff SJ, Worthy TH, Rawlence NJ, Cooper A. 2012 High-resolution coproecology: using coprolites to reconstruct the habits and habitats of New Zealand's extinct upland moa (*Megalapteryx didinus*). *PLoS ONE* **7**, e40025. (doi:10.1371/journal.pone.0040025)
91. Wood JR, Wilmshurst JM, Rawlence NJ, Bonner KI, Worthy TH, Kinsella JM, Cooper A. 2013 A megafauna's microfauna: gastrointestinal parasites of New Zealand's extinct moa (Aves: Dinornithiformes). *PLoS ONE* **8**, e57315. (doi:10.1371/journal.pone.0057315)
92. Brunton-Martin A, Wood JR, Gaskett AC. 2024 Evidence for adaptation of colourful truffle-like fungi for birds in Aotearoa-New Zealand. *Sci. Rep.* **14**, 18908. (doi:10.1038/s41598-024-67333-x)
93. Boast AP, Wood JR, Bolstridge N, Perry GLW, Wilmshurst JM. 2023 Ancient and modern scats record broken ecological interactions and a decline in dietary breadth of the critically endangered kākāpō parrot (*Strigops habroptilus*). *Front. Ecol. Evol.* **11**, 157. (doi:10.3389/fevo.2023.1058130)



94. White TJ, Bruns T, Lee S, Taylor J. 1990 Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR protocols: a guide to methods and applications* (eds M Innis, DH Gelfand, JJ Sninsky, TJ White), pp. 315–322. New York, NY: Academic Press. (doi:10.1016/B978-0-12-372180-8.50042-1)
95. Epp LS *et al.* 2012 New environmental metabarcodes for analysing soil DNA: potential for studying past and present ecosystems. *Mol. Ecol.* **21**, 1821–1833. (doi:10.1111/j.1365-294X.2012.05537.x)
96. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009 BLAST+: architecture and applications. *BMC Bioinform.* **10**, 421. (doi:10.1186/1471-2105-10-421)
97. Huson DH, Beier S, Flade I, Górski A, El-Hadidi M, Mitra S, Ruscheweyh HJ, Tappu R. 2016 MEGAN Community Edition: interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS Comput. Biol.* **12**, e1004957. (doi:10.1371/journal.pcbi.1004957)
98. Darriba D, Taboada GL, Doallo R, Posada D. 2012 jModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods.* **9**, 772. (doi:10.1038/nmeth.2109)
99. Boast AP, Wood C JR, Bolstridge J, Perry N, Wilmshurst GLW. 2024 Supporting data for the manuscript 'DNA and spores from coprolites reveal that colourful truffle-like fungi endemic to NZ were consumed by extinct moa (Dinornithiformes)'. DataStore. (doi:10.7931/8z32-a212)
100. Coluccio AE, Rodriguez RK, Kernan MJ, Neiman AM. 2008 The yeast spore wall enables spores to survive passage through the digestive tract of *Drosophila*. *PLoS ONE* **3**, e2873. (doi:10.1371/journal.pone.0002873)
101. Wood J, Richardson S, McGlone M, Wilmshurst J. 2020 The diets of moa (Aves: Dinornithiformes). *N. Z. J. Ecol.* **44**, 3397. (doi:10.20417/nzjcol.44.3)
102. Simpson J. 1998 Why don't more birds eat more fungi. *Aust. Mycol. Newsl.* **17**, 67–68. <https://www.australasianmycologicalsociety.com/volume-17-3>
103. Crome FHJ. 1976 Some observations on the biology of the cassowary in northern Queensland. *Emu* **76**, 8–14. (doi:10.1071/mu9760008)
104. Wood JR, Vermeulen MJ, Bolstridge N, Briden S, Cole TL, Rivera-Perez J, Shepherd LD, Rawlence NJ, Wilmshurst JM. 2021 Mid-Holocene coprolites from southern New Zealand provide new insights into the diet and ecology of the extinct little bush moa (*Anomalopteryx didiformis*). *Quat. Sci. Rev.* **263**, 106992. (doi:10.1016/j.quascirev.2021.106992)
105. Carpenter JK, Wood JR, Wilmshurst JM, Kelly D. 2018 An avian seed dispersal paradox: New Zealand's extinct megafaunal birds did not disperse large seeds. *Proc. R. Soc. B* **285**, 20180352. (doi:10.1098/rspb.2018.0352)
106. Perry GLW, Wilmshurst JM, Wood JR. 2021 Reconstructing ecological functions provided by extinct fauna using allometrically informed simulation models: An in silico framework for 'movement palaeoecology'. *Funct. Ecol.* **35**, 2575–2592. (doi:10.1111/1365-2435.13904)
107. Worthy T, Scofield R. 2012 Twenty-first century advances in knowledge of the biology of moa (Aves: Dinornithiformes): a new morphological analysis and moa diagnoses revised. *N. Z. J. Zool.* **39**, 87–153. (doi:10.1080/03014223.2012.665060)
108. Frei S, Ortmann S, Reutlinger C, Kreuzer M, Hatt JM, Clauss M. 2015 Comparative digesta retention patterns in ratites. *Auk*. **132**, 119–131. (doi:10.1642/auk-14-144.1)
109. Bunce M *et al.* 2009 The evolutionary history of the extinct ratite moa and New Zealand Neogene paleogeography. *Proc. Natl Acad. Sci. USA* **106**, 20646–20651. (doi:10.1073/pnas.0906660106)
110. Pierre JP. 1999 Reintroduction of the South Island saddleback (*Philesturnus carunculatus carunculatus*): dispersal, social organisation and survival. *Biol. Conserv.* **89**, 153–159. (doi:10.1016/S0006-3207(98)00139-6)
111. Menzies IJ, Burns K. 2010 Temporal shifts in the pair-bond dynamics of New Zealand robins (*Petroica australis*). *N. Z. J. Ecol.* 265–268.
112. Godínez-Alvarez H, Ríos-Casanova L, Peco B. 2020 Are large frugivorous birds better seed dispersers than medium- and small-sized ones? Effect of body mass on seed dispersal effectiveness. *Ecol. Evol.* **10**, 6136–6143. (doi:10.1002/ece3.6285)
113. Carpenter JK, Innes JG, Wood JR, Lyver PO. 2021 Good predators: the roles of weka (*Gallirallus australis*) in New Zealand's past and present ecosystems. *N.Z.J.Ecol.* **45**. (doi:10.20417/nzjcol.45.15)
114. Checklist Committee (OSNZ). 2022 Checklist of the Birds of New Zealand. In *Ornithological society of new zealand occasional publication no. 1*, 5th edn. Wellington: Ornithological Society of New Zealand.
115. Lebel T, Davoodian N, Bloomfield M, Syme K, May T, Hosaka K, Castellano M. 2022 A mixed bag of sequester fungi from five different families: Boletaceae, Russulaceae, Psathyrellaceae, Strophariaceae, and Hysterangiaceae. *Swainsona* **36**, 33–65. <https://www.jstor.org/stable/27284114>
116. Cribb J. 1994 The fungus *Secotium fragariosum* newly recorded from Lord Howe Island. *Qld. Nat.* **33**, 32.
117. Grandcolas P, Muriene J, Robillard T, Desutter-Grandcolas L, Jourdan H, Guilbert E, Deharveng L. 2008 New Caledonia: a very old Darwinian island? *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* **363**, 3309–3317. (doi:10.1098/rstb.2008.0122)
118. Hutton I, Parkes JP, Sinclair ARE. 2007 Reassembling island ecosystems: the case of Lord Howe Island. *Anim. Conserv.* **10**, 22–29. (doi:10.1111/j.1469-1795.2006.00077.x)
119. Gasparini B. 2014 Cortinari (Agaricales) revised taxonomy: new species names or combinations. *Mycosphere* **5**, 541–544. (doi:10.5943/mycosphere/5/4/6)
120. Wood JR, Dickie IA, Moeller HV, Peltzer DA, Bonner KI, Rattray G, Wilmshurst JM. 2015 Novel interactions between non-native mammals and fungi facilitate establishment of invasive pines. *J. Ecol.* **103**, 121–129. (doi:10.1111/1365-2745.12345)
121. Claridge AW, May TW. 1994 Mycophagy among Australian mammals. *Aust. J. Ecol.* **19**, 251–275. (doi:10.1111/j.1442-9993.1994.tb00489.x)
122. Dickie IA, Davis M, Carswell FE. 2012 Quantification of mycorrhizal limitation in beech spread. *N. Z. J. Ecol.* 210–215. <https://newzealandecology.org/nzje/3032>
123. Baylis G. 1980 Mycorrhizas and the spread of beech. *N.Z.J.Ecol.* **3**, 151–153. <https://newzealandecology.org/nzje/1485>
124. Aguirre F, Nouhra E, Urcelay C. 2021 Native and non-native mammals disperse exotic ectomycorrhizal fungi at long distances from pine plantations. *Fungal Ecol.* **49**, 101012. (doi:10.1016/j.funeco.2020.101012)
125. Soteras F, Ibarra C, Geml J, Barrios-García MN, Domínguez LS, Nouhra ER. 2017 Mycophagy by invasive wild boar (*Sus scrofa*) facilitates dispersal of native and introduced mycorrhizal fungi in Patagonia, Argentina. *Fungal Ecol.* **26**, 51–58. (doi:10.1016/j.funeco.2016.11.008)
126. Urcelay C, Longo S, Geml J, Tecco PA, Nouhra E. 2017 Co-invasive exotic pines and their ectomycorrhizal symbionts show capabilities for wide distance and altitudinal range expansion. *Fungal Ecol.* **25**, 50–58. (doi:10.1016/j.funeco.2016.11.002)
127. Boast AP, Wood J, Cooper J, Boldstridge N, Perry GLW, Wilmshurst JM. 2024 Supplementary material from: DNA and spores from coprolites reveal that colourful truffle-like fungi endemic to NZ were consumed by extinct moa (Dinornithiformes). Figshare. (doi:10.6084/m9.figshare.c.7584142)